

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use FAMCICLOVIR TABLETS safely and effectively. See full prescribing information for FAMCICLOVIR TABLETS.

### FAMCICLOVIR tablets, for oral use

Initial U.S. Approval: 1994

### INDICATIONS AND USAGE

Famciclovir, a prodrug of penciclovir, is a deoxy nucleoside analog DNA polymerase inhibitor indicated for:

#### Immunocompetent Adult Patients (1.1)

- Herpes labialis (cold sores)
  - Treatment of recurrent episodes
- Genital herpes
  - Treatment of recurrent episodes
  - Suppressive therapy of recurrent episodes
- Herpes zoster (shingles)

#### Human Immunodeficiency Virus (HIV)-Infected Adult Patients (1.2)

- Treatment of recurrent episodes of orolabial or genital herpes

#### Limitation of Use

The efficacy and safety of famciclovir have not been established for:

- Patients with first episode of genital herpes
- Patients with ophthalmic zoster
- Immunocompromised patients other than for the treatment of recurrent episodes of orolabial or genital herpes in HIV-infected patients
- Black and African American patients with recurrent genital herpes

### DOSAGE AND ADMINISTRATION

| Immunocompetent Adult Patients (2.1) |           |             |  |
|--------------------------------------|-----------|-------------|--|
| Herpes labialis (cold sores)         | labialis  | (cold)      | 1500 mg as a single dose   |
| Genital herpes                       | recurrent | Suppressive | 1000 mg twice daily for episodes<br>1 day 250 mg twice daily therapy |

### FULL PRESCRIBING INFORMATION: CONTENTS\*

|                                      |  |
|--------------------------------------|--|
| <b>1 INDICATIONS AND USAGE</b>       | 1.1 Immunocompetent Adult Patients<br>1.2 HIV-Infected Adult Patients  |
| <b>2 DOSAGE AND ADMINISTRATION</b>   | 2.1 Dosing Recommendation in Immunocompetent Adult Patients<br>2.2 Dosing Recommendation in HIV-Infected Adult Patients<br>2.3 Dosing Recommendation in Patients with Renal Impairment   |
| <b>3 DOSAGE FORMS AND STRENGTHS</b>  |  |
| <b>4 CONTRAINDICATIONS</b>           |  |
| <b>5 WARNINGS AND PRECAUTIONS</b>    | 5.1 Acute Renal Failure  |
| <b>6 ADVERSE REACTIONS</b>           | 6.1 Clinical Trials Experience in Adult Patients<br>6.2 Postmarketing Experience   |
| <b>7 DRUG INTERACTIONS</b>           | 7.1 Potential for Famciclovir Tablets to Affect Other Drugs<br>7.2 Potential for Other Drugs to Affect Penciclovir   |
| <b>8 USE IN SPECIFIC POPULATIONS</b> | 8.1 Pregnancy<br>8.2 Lactation<br>8.3 Females and Males of Reproductive Potential<br>8.4 Pediatric Use<br>8.5 Geriatric Use<br>8.6 Patients with Renal Impairment<br>8.7 Patients with Hepatic Impairment<br>8.8 Black and African American Patients |

### FULL PRESCRIBING INFORMATION

#### 1 INDICATIONS AND USAGE

##### 1.1 Immunocompetent Adult Patients

Herpes labialis (cold sores): Famciclovir tablets are indicated for the treatment of recurrent herpes labialis in adult patients.

##### Genital herpes:

Recurrent episodes: Famciclovir tablets are indicated for the treatment of recurrent episodes of genital herpes. The efficacy of famciclovir tablets when initiated more than 6 hours after onset of symptoms or lesions has not been established.

Suppressive therapy: Famciclovir tablets are indicated for chronic suppressive therapy of recurrent episodes of genital herpes in adult patients. The efficacy and safety of famciclovir tablets for the suppression of recurrent genital herpes beyond 1 year have not been established.

Herpes zoster (shingles): Famciclovir tablets are indicated for the treatment of herpes zoster in adult patients. The efficacy of famciclovir tablets when initiated more than 72 hours after onset of rash has not been established.

##### 1.2 HIV-Infected Adult Patients

Recurrent orolabial or genital herpes: Famciclovir tablets are indicated for the treatment of recurrent episodes of orolabial or genital herpes in HIV-infected adults. The efficacy of famciclovir tablets when initiated more than 48 hours after onset of symptoms or lesions has not been established.

|   |                                 |
|---|---------------------------------|
| Herpes zoster (shingles)                          | 500 mg every 8 hours for 7 days |
| HIV-Infected Adult Patients (2.2)                 |                                 |
| Recurrent episodes of orolabial or genital herpes | 500 mg twice daily for 7 days   |

Patients with renal impairment: Adjust dose based on creatinine clearance. (2.3)

### DOSAGE FORMS AND STRENGTHS

Tablets: 125 mg, 250 mg, 500 mg (3)

### CONTRAINDICATIONS

Known hypersensitivity to the product, its components, or Denavir® (penciclovir cream). (4)

### WARNINGS AND PRECAUTIONS

Acute renal failure: May occur in patients with underlying renal disease who receive higher than recommended doses of famciclovir tablets for their level of renal function. Reduce dosage in patients with renal impairment. (2.3, 8.6)

### ADVERSE REACTIONS

The most common adverse events reported in at least 1 indication by greater than 10% of adult patients are headache and nausea. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Cipla Limited, at 1-866-604-3268 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

### DRUG INTERACTIONS

Probenecid: May increase penciclovir levels. Monitor for evidence of penciclovir toxicity. (7.2)

### See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 1/2020

#### Limitation of Use

The efficacy and safety of famciclovir tablets have not been established for:

- Patients with first episode of genital herpes
- Patients with ophthalmic zoster
- Immunocompromised patients other than for the treatment of recurrent orolabial or genital herpes in HIV-infected patients
- Black and African American patients with recurrent genital herpes

### DOSAGE AND ADMINISTRATION

Famciclovir tablets may be taken with or without food.

#### 2.1 Dosing Recommendation in Immunocompetent Adult Patients

Herpes labialis (cold sores): The recommended dosage of famciclovir tablets for the treatment of recurrent herpes labialis is 1500 mg as a single dose. Therapy should be initiated at the first sign or symptom of herpes labialis (e.g., tingling, itching, burning, pain, or lesion).

##### Genital herpes:

Recurrent episodes: The recommended dosage of famciclovir tablets for the treatment of recurrent episodes of genital herpes is 1000 mg twice daily for 1 day. Therapy should be initiated at the first sign or symptom of a recurrent episode (e.g., tingling, itching, burning, pain, or lesion).

Suppressive therapy: The recommended dosage of famciclovir tablets for chronic suppressive therapy of recurrent episodes of genital herpes is 250 mg twice daily.

Herpes zoster (shingles): The recommended dosage of famciclovir tablets for the treatment of herpes zoster is 500 mg every 8 hours for 7 days. Therapy should be initiated as soon as herpes zoster is diagnosed.

#### 2.2 Dosing Recommendation in HIV-Infected Adult Patients

Recurrent orolabial or genital herpes: The recommended dosage of famciclovir tablets for the treatment of recurrent orolabial or genital herpes in HIV-infected patients is 500 mg twice daily for 7 days. Therapy should be initiated at the first sign or symptom of a recurrent episode (e.g., tingling, itching, burning, pain, or lesion).

#### 2.3 Dosing Recommendation in Patients with Renal Impairment

Dosage recommendations for adult patients with renal impairment are provided in Table 1 (see *Use in Specific Populations (8.6)*, *Clinical Pharmacology (12.3)*).

#### Table 1: Dosage Recommendations for Adult Patients with Renal Impairment

| Indication and Normal Dosage Regimen   | Creatinine Clearance (mL/min) | Adjusted Dosage Regimen Dose (mg) | Dosing Interval                |
|--|-------------------------------|-----------------------------------|--------------------------------|
| <b>Single-Day Dosing Regimens</b>  |                               |                                   |                                |
| Recurrent Genital Herpes<br>1000 mg every 12 hours for 1 day                         | ≥ 60                          | 1000                              | every 12 hours for 1 day       |
|  | 40-59                         | 500                               | every 12 hours for 1 day       |
|  | 20-39                         | 500                               | single dose                    |
|  | < 20                          | 250                               | single dose                    |
|  | HD*                           | 250                               | single dose following dialysis |
| Recurrent Herpes Labialis<br>1500 mg single dose                                     | ≥ 60                          | 1500                              | single dose                    |
|  | 40-59                         | 750                               | single dose                    |
|  | 20-39                         | 500                               | single dose                    |
|  | < 20                          | 250                               | single dose                    |
|  | HD*                           | 250                               | single dose following dialysis |
| <b>Multiple-Day Dosing Regimens</b>  |                               |                                   |                                |
| Herpes Zoster<br>500 mg every 8 hours  | ≥ 60                          | 500                               | every 8 hours                  |
|  | 40-59                         | 500                               | every 12 hours                 |
|  | 20-39                         | 500                               | every 24 hours                 |
|  | < 20                          | 250                               | every 24 hours                 |
|  | HD*                           | 250                               | following each dialysis        |
|  | ≥ 40                          | 250                               | every 12 hours                 |
| Suppression of Recurrent Genital Herpes<br>250 mg every 12 hours                     | ≥ 60                          | 125                               | every 12 hours                 |
|  | < 20                          | 125                               | every 24 hours                 |
|  | HD*                           | 125                               | following each dialysis        |
|  | ≥ 40                          | 500                               | every 12 hours                 |
| Recurrent Orolabial or Genital Herpes in HIV-Infected Patients 500 mg every 12 hours | ≥ 60                          | 500                               | every 24 hours                 |
|  | < 20                          | 250                               | every 24 hours                 |
|  | HD*                           | 250                               | following each dialysis        |

##### \*Hemodialysis

#### 3 DOSAGE FORMS AND STRENGTHS

Famciclovir tablets are available in 3 strengths:

- 125 mg: White coloured, circular, film-coated, biconvex tablet debossed with 'C' and '269' on one side and plain on other side.
- 250 mg: White coloured, circular, film-coated, biconvex tablet debossed with 'C' and '271' on one side and plain on other side.
- 500 mg: White coloured, capsule shaped, film-coated, biconvex tablets debossed with 'C272' on one side and plain on other side.

#### 4 CONTRAINDICATIONS

Famciclovir tablets are contraindicated in patients with known hypersensitivity to the product, its components, or Denavir® (penciclovir cream).

#### 5 WARNINGS AND PRECAUTIONS

##### 5.1 Acute Renal Failure

Cases of acute renal failure have been reported in patients with underlying renal disease who have received inappropriately high doses of famciclovir for their level of renal function. Dosage reduction is recommended when administering famciclovir to patients with renal impairment (see *Dosage and Administration (2.3)*, *Use in Specific Populations (8.6)*).

#### 6 ADVERSE REACTIONS

Acute renal failure is discussed in greater detail in other sections of the label (see *Warnings and Precautions (5)*).

The most common adverse events reported in at least 1 indication by greater than 10% of adult patients treated with famciclovir are headache and nausea.

#### 6.1 Clinical Trials Experience in Adult Patients

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Immunocompetent patients: The safety of famciclovir has been evaluated in active-and placebo-controlled clinical studies involving 816 famciclovir-treated patients with herpes zoster (famciclovir, 250 mg three times daily to 750 mg three times daily); 163 famciclovir-treated patients with recurrent genital herpes (famciclovir, 1000 mg twice daily); 1,197 patients with recurrent genital herpes treated with famciclovir as suppressive therapy (125 mg once daily to 250 mg three times daily) of which 570 patients received famciclovir (open-labeled and/or double-blind) for at least 10 months; and 447 famciclovir-treated patients with herpes labialis (famciclovir, 1500 mg once daily or 750 mg twice daily). Table 2 lists selected adverse events.

Table 2: Selected Adverse Events (all grades and without regard to causality) Reported by Greater Than or Equal to 2% of Patients in Placebo-Controlled Famciclovir Trials

| Events                       | Incidence                  |                   |                                       |                   |   |                   |                              |                   |
|------------------------------|----------------------------|-------------------|---------------------------------------|-------------------|---|-------------------|------------------------------|-------------------|
|                              | Herpes Zoster <sup>1</sup> |                   | Recurrent Genital Herpes <sup>2</sup> |                   | Genital Herpes-Suppression <sup>3</sup> |                   | Herpes Labialis <sup>4</sup> |                   |
|                              | Famciclovir (n=273) %      | Placebo (n=146) % | Famciclovir (n=163) %                 | Placebo (n=166) % | Famciclovir (n=458) %                   | Placebo (n=463) % | Famciclovir (n=447) %        | Placebo (n=254) % |
| <b>Nervous System</b>        |                            |                   |                                       |                   |   |                   |                              |                   |
| Headache                     | 22.7                       | 17.8              | 13.5                                  | 5.4               | 39.3                                    | 42.9              | 8.5                          | 6.7               |
| Paresthesia                  | 2.6                        | 0.0               | 0.0                                   | 0.0               | 0.9                                     | 0.0               | 0.0                          | 0.0               |
| Migraine                     | 0.7                        | 0.7               | 0.6                                   | 0.6               | 3.1                                     | 0.0               | 0.2                          | 0.0               |
| <b>Gastrointestinal</b>      |                            |                   |                                       |                   |   |                   |                              |                   |
| Nausea                       | 12.5                       | 11.6              | 2.5                                   | 3.6               | 7.2                                     | 9.5               | 2.2                          | 3.9               |
| Diarrhea                     | 7.7                        | 4.8               | 4.9                                   | 1.2               | 9.0                                     | 9.5               | 1.6                          | 0.8               |
| Vomiting                     | 4.8                        | 3.4               | 1.2                                   | 0.6               | 3.1                                     | 1.6               | 0.7                          | 0.0               |
| Flatulence                   | 1.5                        | 0.7               | 0.6                                   | 0.0               | 4.8                                     | 1.6               | 0.2                          | 0.0               |
| Abdominal Pain               | 1.1                        | 3.4               | 0.0                                   | 1.2               | 7.9                                     | 7.9               | 0.2                          | 0.4               |
| <b>Body as a Whole</b>       |                            |                   |                                       |                   |   |                   |                              |                   |
| Fatigue                      | 4.4                        | 3.4               | 0.6                                   | 0.0               | 4.8                                     | 3.2               | 1.6                          | 0.4               |
| <b>Skin and Appendages</b>   |                            |                   |                                       |                   |   |                   |                              |                   |
| Pruritus                     | 3.7                        | 2.7               | 0.0                                   | 0.6               | 2.2                                     | 0.0               | 0.0                          | 0.0               |
| Rash                         | 0.4                        | 0.7               | 0.0                                   | 0.0               | 3.3                                     | 1.6               | 0.0                          | 0.0               |
| <b>Reproductive (Female)</b> |                            |                   |                                       |                   |   |                   |                              |                   |
| Dysmenorrhea                 | 0.0                        | 0.7               | 1.8                                   | 0.6               | 7.6                                     | 6.3               | 0.4                          | 0.0               |

<sup>1</sup>Patients may have entered into more than one clinical trial.

<sup>2</sup>7 days of treatment

<sup>3</sup>1 day of treatment

<sup>4</sup>daily treatment

Table 3 lists selected laboratory abnormalities in genital herpes suppression trials.

#### Table 3: Selected Laboratory Abnormalities in Genital Herpes Suppression Studies<sup>1</sup>

| Parameter                     | Famciclovir (n=660) <sup>1</sup> % | Placebo (n=210) <sup>1</sup> % |
|-------------------------------|------------------------------------|--------------------------------|
| Anemia (<0.8 x NRL)           | 0.1                                | 0.0                            |
| Leukopenia (<0.75 x NRL)      | 1.3                                | 0.9                            |
| Neutropenia (<0.8 x NRL)      | 3.2                                | 1.5                            |
| AST (SGOT) (>2 x NRH)         | 2.3                                | 1.2                            |
| ALT (SGPT) (>2 x NRH)         | 3.2                                | 1.5                            |
| Total Bilirubin (>1.5 x NRH)  | 1.9                                | 1.2                            |
| Serum Creatinine (>1.5 x NRH) | 0.2                                | 0.3                            |
| Amylase (>1.5 x NRH)          | 1.5                                | 1.9                            |
| Lipase (>1.5 x NRH)           | 4.9                                | 4.7                            |

<sup>1</sup>Percentage of patients with laboratory abnormalities that were increased or decreased from baseline and were outside of specified ranges.

<sup>2</sup>n values represent the minimum number of patients assessed for each laboratory parameter.

NRH=Normal Range High.

NRL=Normal Range Low.

HIV-infected patients: In HIV-infected patients, the most frequently reported adverse events for famciclovir (500 mg twice daily; n=150) and acyclovir (400 mg, 5x/day; n=143), respectively, were headache (17% vs. 15%), nausea (11% vs. 13%), diarrhea (7% vs. 11%), vomiting (5% vs. 4%), fatigue (4% vs. 2%), and abdominal pain (3% vs. 6%).

#### 6.2 Postmarketing Experience

The adverse events listed below have been reported during post-approval use of famciclovir. Because these events are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure:

*Blood and lymphatic system disorders:* Thrombocytopenia

*Hepatobiliary disorders:* Abnormal liver function tests, cholestatic jaundice

*Immune system disorders:* Anaphylactic shock, anaphylactic reaction

*Nervous system disorders:* Dizziness, somnolence, seizure

*Psychiatric disorders:* Confusion (including delirium, disorientation, and confusional state occurring predominantly in the elderly), hallucinations

*Skin and subcutaneous tissue disorders:* Urticaria, erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, angioedema (e.g., face, eyelid, periorbital, and pharyngeal edema), hypersensitivity vasculitis

*Cardiac disorders:* Palpitations

#### 7 DRUG INTERACTIONS

##### 7.1 Potential for Famciclovir Tablets to Affect Other Drugs

The steady-state pharmacokinetics of digoxin were not altered by concomitant administration of multiple doses of famciclovir (500 mg three times daily). No clinically significant effect on the pharmacokinetics of zidovudine, its metabolite zidovudine glucuronide, or emtricitabine was observed following a single oral dose of 500 mg famciclovir coadministered with zidovudine or emtricitabine.

An *in vitro* study using human liver microsomes suggests that famciclovir is not an inhibitor of CYP3A4 enzymes.

##### 7.2 Potential for Other Drugs to Affect Famciclovir

No clinically significant alterations in famciclovir pharmacokinetics were observed following single-dose administration of 500 mg famciclovir after pretreatment with multiple doses of allopurinol, cimetidine, theophylline, zidovudine, promethazine, when given shortly after an antacid (magnesium and aluminum hydroxide), or concomitantly with emtricitabine. No clinically significant effect on famciclovir pharmacokinetics was observed following multiple-dose (three times daily) administration of famciclovir (500 mg) with multiple doses of digoxin.

Concurrent use with probenecid or other drugs significantly eliminated by active renal tubular secretion may result in increased plasma concentrations of famciclovir.

The conversion of 6-deoxy famciclovir to famciclovir is catalyzed by aldehyde oxidase. Interactions with other drugs metabolized by this enzyme and/or inhibiting this enzyme could potentially occur. Clinical interaction studies of famciclovir with cimetidine and promethazine, *in vitro* inhibitors of aldehyde oxidase, did not show relevant effects on the formation of famciclovir. Raloxifene, a potent aldehyde oxidase inhibitor *in vitro*, could decrease the formation of famciclovir. However, a clinical drug-drug interaction study to determine the magnitude of interaction between famciclovir and raloxifene has not been conducted.

#### 8 USE IN SPECIFIC POPULATIONS

##### 8.1 Pregnancy

###### Risk Summary

Available data from pharmacovigilance reports with famciclovir use in pregnant women have not identified a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. There are risks to the fetus associated with untreated herpes simplex virus during pregnancy (see *Clinical Considerations*). After oral administration, famciclovir (prodrug) is converted to penciclovir (active drug). In animal reproduction studies with famciclovir, no evidence of adverse developmental outcomes was observed at systemic exposures of famciclovir (AUC) slightly higher than those at the maximum recommended human dose (MRHD) of famciclovir (see *Data*).

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage is clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Healthcare providers are encouraged to report pregnancies and pregnancy outcomes to Cipla Limited, at 1-866-604-3268

#### Clinical Considerations

##### Disease-associated maternal and/or embryo-fetal risk

The risk of neonatal herpes infection varies from 30% to 50% for genital herpes simplex virus (HSV) infections that occur in late pregnancy (third trimester), whereas in early pregnancy, infection carries a risk of about 1%. A primary herpes outbreak during the first trimester of pregnancy has been associated with neonatal chorioretinitis, microcephaly and, in rare cases, skin lesions. In very rare cases, transplacental transmission can occur resulting in congenital infection, including microcephaly, hepatosplenomegaly, intrauterine growth restriction and stillbirth. Co-infection with HSV increases the risk of perinatal HIV transmission in women who had a clinical diagnosis of genital herpes during pregnancy.

##### Data

###### Animal Data

Famciclovir was administered orally to pregnant rats and rabbits (up to 1000 mg/kg/day) on gestation Day(s) 6 to 15, and to rats on gestation Day 15 to lactation/post-partum Day 25. No adverse effects on embryo-fetal (rats and rabbits) or pre/post-natal (rats) development were observed up to the highest dose tested. During organogenesis, systemic exposures of famciclovir (active metabolite) were 3.4 times (rats) and 1.6 times (rabbits) the human systemic exposure of famciclovir based on AUC at the MRHD.

##### 8.2 Lactation

There are no data on the presence of famciclovir (prodrug) or penciclovir (active drug) in human milk, the effects on the breastfed infant, or the effects on milk production. Animal data indicate that famciclovir is present in the milk of lactating rats (see *Data*). The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for famciclovir and any potential adverse effects on the breastfed infant from famciclovir or from the underlying maternal condition.

##### Data

Penciclovir was the primary drug-related component excreted into the milk of lactating rats following a single oral dose of 40 mg/kg on lactation Day 12, with milk concentrations of up to approximately 8 times that of maternal plasma concentrations observed 0.5 hours postdose.

##### 8.3 Females and Males of Reproductive Potential

###### Fertility

Decreased fertility, due to testicular toxicity, was observed in male animals following repeated administration of famciclovir or penciclovir (see *Nonclinical Toxicology (13.1)*). In two placebo-controlled studies, 130 men with a history of recurrent genital herpes received either oral famciclovir (250 mg twice daily; n=66) or placebo (n=64) therapy for 18 weeks. The men were otherwise healthy and had a normal sperm profile prior to treatment. There was no evidence of significant effects on sperm count, motility or morphology during famciclovir treatment or during an 8-week follow-up.

##### 8.4 Pediatric Use

The efficacy of famciclovir have not been established in pediatric patients. The pharmacokinetic profile and safety of famciclovir (experimental granules mixed with OraSweet® or tablets) were studied in 3 open-label studies.

Study 1 was a single-dose pharmacokinetic and safety study in infants 1 to less than 1 year of age who had an active herpes simplex virus (HSV) infection or who were at risk for HSV infection. Eighteen subjects were enrolled and received a single dose of famciclovir experimental granules mixed with OraSweet based on the patient's body weight (doses ranged from 25 mg to 175 mg). These doses were selected to provide famciclovir systemic exposures similar to the famciclovir systemic exposures observed in adults after administration of 500 mg famciclovir. The efficacy and safety of famciclovir have not been established as suppressive therapy in infants following neonatal HSV infections. In addition, the efficacy cannot be extrapolated from adults to infants because there is no similar disease in adults. Therefore, famciclovir is not recommended in infants.

Study 2 was an open-label, single-dose pharmacokinetic, multiple-dose safety study of famciclovir experimental granules mixed with OraSweet in children 1 to less than 12 years of age with clinically suspected HSV or varicella zoster virus (VZV) infection. Fifty-one subjects were enrolled in the pharmacokinetic part of the study and received a single body weight adjusted dose of famciclovir (doses ranged from 125 mg to 500 mg). These doses were selected to provide famciclovir systemic exposures similar to the famciclovir systemic exposures observed in adults after administration of 500 mg famciclovir. Based on the pharmacokinetic data observed with these doses in children, a new weight-based dosing algorithm was designed and used in the multiple-dose safety part of the study. Pharmacokinetic data were not obtained with the revised weight-based dosing algorithm.

A total of 100 patients were enrolled in the multiple-dose safety part of the study; 47 subjects with active or latent HSV infection and 53 subjects with chickenpox. Patients with active or latent HSV infection received famciclovir twice a day for 7 days. The daily dose of famciclovir ranged from 150 mg to 500 mg twice daily depending on the patient's body weight. Patients with chickenpox received famciclovir three times daily for 7 days. The daily dose of famciclovir ranged from 150 mg to 500 mg three times daily depending on the patient's body weight. The clinical adverse events and laboratory test abnormalities observed in this study were similar to those seen in adults. The available data are insufficient to support the use of famciclovir for the treatment of children 1 to less than 12 years of age with chickenpox or infections due to HSV for the following reasons:

*Chickenpox:* The efficacy of famciclovir for the treatment of chickenpox has not been established in either pediatric or adult



### What are the possible side effects of famciclovir tablets?

The most common side effects of famciclovir tablets are headache and nausea.

These are not all the possible side effects of famciclovir tablets.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### How should I store famciclovir tablets?

- Store famciclovir tablets at 20°C to 25°C (68°F to 77°F) [see USP Controlled Room Temperature].

### Keep famciclovir tablets and all medicines out of reach from children.

### General information about the safe and effective use of famciclovir tablets

Famciclovir tablets are not a cure for cold sores or genital herpes. It is not known if famciclovir tablets can stop the spread of herpes to others. If you are sexually active, you can pass herpes to your partner even if you are taking famciclovir tablets. Herpes can be passed to your partner even if you do not have active symptoms. You should continue to practice safer sex to lower the chances of spreading genital herpes to others. Do not have sexual contact with your partner during an outbreak of genital herpes or if you have any symptoms of genital herpes. Ask your healthcare provider for more information about safer sex practices.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use famciclovir tablets for a condition for which it was not prescribed. Do not give famciclovir tablets to other people, even if they have the same symptoms you have. It may harm them. You can ask your healthcare provider or pharmacist for information about famciclovir tablets that is written for health professionals.

### What are the ingredients in famciclovir tablets?

#### Active ingredient: famciclovir

**Inactive ingredients:** croscarmellose sodium, crospovidone, hypromellose 2910, lactose monohydrate, magnesium stearate, polyethylene glycols 6000 and titanium dioxide.

Disclaimer: Other Brands listed are the registered trademarks of their respective owners and are not trademarks of Cipla Limited.

Manufactured by: Cipla Ltd., Verna Goa, India

Manufactured for: Cipla USA, Inc.

10 Independence Boulevard, Suite 300

Warren, NJ 07059

Revised: 1/2020

In a multiple-dose study of famciclovir conducted in subjects with varying degrees of renal impairment (n=18), the pharmacokinetics of penciclovir were comparable to those after single doses.

A dosage adjustment is recommended for patients with renal impairment [see *Dosage and Administration* (2.3)].

#### 8.7 Patients with Hepatic Impairment

Mild or moderate hepatic impairment (chronic hepatitis [n=6], chronic ethanol abuse [n=8], or primary biliary cirrhosis [n=1]) had no effect on the extent of availability (AUC) of penciclovir following a single dose of 500 mg famciclovir. However, there was a 44% decrease in penciclovir mean maximum plasma concentration (C<sub>max</sub>) and the time to maximum plasma concentration (t<sub>max</sub>) was increased by 0.75 hours in patients with hepatic impairment compared to normal volunteers. No dosage adjustment is recommended for patients with mild or moderate hepatic impairment. The pharmacokinetics of penciclovir has not been evaluated in patients with severe hepatic impairment. Conversion of famciclovir to the active metabolite penciclovir may be impaired in these patients resulting in a lower penciclovir plasma concentrations, and thus possibly a decrease of efficacy of famciclovir [see *Clinical Pharmacology* (12)].

#### 8.8 Black and African American Patients


In a randomized, double-blind, placebo-controlled trial conducted in 304 immunocompetent black and African American adults with recurrent genital herpes there was no difference in median time to healing between patients receiving famciclovir or placebo. In general, the adverse reaction profile was similar to that observed in other famciclovir clinical trials for adult patients [see *Adverse Reactions* (6.1)]. The relevance of these study results to other indications in black and African American patients is unknown [see *Clinical Studies* (14.2)].

#### 10 OVERDOSAGE

Appropriate symptomatic and supportive therapy should be given. Penciclovir is removed by hemodialysis.

#### 11 DESCRIPTION

The active ingredient in famciclovir tablets is famciclovir, an orally administered prodrug of the antiviral agent penciclovir. Chemically, famciclovir is known as 2-[2-(2-amino-9H-purin-9-yl)ethyl]-1,3-propanediol diacetate. Its molecular formula is C<sub>14</sub>H<sub>16</sub>N<sub>6</sub>O<sub>6</sub>; its molecular weight is 321.3. It is a synthetic acyclic guanine derivative and has the following structure



Famciclovir is a white to pale yellow solid. It is freely soluble in acetone and methanol, and sparingly soluble in ethanol and isopropanol. At 25°C famciclovir is freely soluble (greater than 25% w/v) in water initially, but rapidly precipitates as the sparingly soluble (2%-3% w/v) monohydrate. Famciclovir is not hygroscopic below 85% relative humidity. Partition coefficients are: octanol/water (pH 4.8) P=1.09 and octanol/phosphate buffer (pH 7.4) P=2.08.

Famciclovir tablets contain 125 mg, 250 mg or 500 mg of famciclovir, together with the following inactive ingredients: croscarmellose sodium, crospovidone, hypromellose 2910, lactose monohydrate, magnesium stearate, polyethylene glycols 6000 and titanium dioxide.

Meets USP Dissolution Test 2

#### 12 CLINICAL PHARMACOLOGY

##### 12.1 Mechanism of Action

Famciclovir is an orally administered prodrug of the anti-alpha herpes viral agent penciclovir [see *Microbiology* (12.4)].

##### 12.3 Pharmacokinetics

Famciclovir is the diacetyl 6-deoxy analog of the active antiviral compound penciclovir. Following oral administration famciclovir undergoes rapid and extensive metabolism to penciclovir and little or no famciclovir is detected in plasma or urine. Penciclovir is predominantly eliminated unchanged by the kidney. Therefore, the dose of famciclovir needs to be adjusted in patients with different degrees of renal impairment [see *Dosage and Administration* (2.3)].

##### Pharmacokinetics in Adults

**Absorption and Bioavailability:** The absolute bioavailability of penciclovir is 7 ± 8% as determined following the administration of a 500 mg famciclovir oral dose and a 400 mg penciclovir intravenous dose to 12 healthy male subjects.

Penciclovir concentrations increased in proportion to dose over a famciclovir dose range of 125 mg to 1000 mg administered as a single dose. Table 5 shows the mean pharmacokinetic parameters of penciclovir after single administration of famciclovir to healthy male volunteers.

**Table 5: Mean Pharmacokinetic Parameters of Penciclovir in Healthy Adult Subjects\***

| Dose    | AUC (0-inf) <sup>1</sup><br>(mcg hr/mL) | C <sub>max</sub> <sup>1</sup><br>(mcg/mL) | t <sub>max</sub> <sup>2</sup><br>(h) |
|---------|---|---|--------------------------------------|
| 125 mg  | 2.24                                    | 0.8                                       | 0.9                                  |
| 250 mg  | 4.48                                    | 1.6                                       | 0.9                                  |
| 500 mg  | 8.95                                    | 3.3                                       | 0.9                                  |
| 1000 mg | 17.9                                    | 6.6                                       | 0.9                                  |

\* Based on pharmacokinetic data from 17 studies

<sup>1</sup> AUC (0-inf) (mcg hr/mL) = area under the plasma concentration-time profile extrapolated to infinity.

<sup>2</sup> C<sub>max</sub> (mcg/mL) = maximum observed plasma concentration.

t<sub>max</sub> (h) = time to C<sub>max</sub>

Following oral single-dose administration of 500 mg famciclovir to 7 patients with herpes zoster, the AUC (mean ± SD), C<sub>max</sub> and t<sub>max</sub> were 12.1±1.7 mcg hr/mL, 4.0±0.7 mcg/mL, and 0.7±0.2 hours, respectively. The AUC of penciclovir was approximately 35% greater in patients with herpes zoster as compared to healthy volunteers. Some of this difference may be due to differences in renal function between the 2 groups.

There is no accumulation of penciclovir after the administration of 500 mg famciclovir three times daily for 7 days.

Penciclovir C<sub>max</sub> decreased approximately 50% and t<sub>max</sub> was delayed by 1.5 hours when a capsule formulation of famciclovir was administered with food (nutritional content was approximately 910 Kcal and 26% fat). There was no effect on the extent of availability (AUC) of penciclovir. There was an 18% decrease in C<sub>max</sub> and a delay in t<sub>max</sub> of about 1 hour when famciclovir was given 2 hours after a meal as compared to its administration 2 hours before a meal. Because there was no effect on the extent of systemic availability of penciclovir, famciclovir can be taken without regard to meals.

**Distribution:** The volume of distribution (Vdβ) was 1.08±0.17 L/kg in 12 healthy male subjects following a single intravenous dose of penciclovir at 400 mg administered as a 1-hour intravenous infusion. Penciclovir is less than 20% bound to plasma proteins over the concentration range of 0.1 to 20 mcg/mL. The blood/plasma ratio of penciclovir is approximately 1.

**Metabolism:** Following oral administration, famciclovir is deacetylated and oxidized to form penciclovir. Metabolites that are inactive include 6-deoxy penciclovir, monoacetylated penciclovir, and 6-deoxy monoacetylated penciclovir (5%, less than 0.5% and less than 0.5% of the dose in the urine, respectively). Little or no famciclovir is detected in plasma or urine. An *in vitro* study using human liver microsomes demonstrated that cyclooxenone P450 does not play an important role in famciclovir metabolism. The conversion of 6-deoxy penciclovir to penciclovir is catalyzed by aldehyde oxidase. Simeidine and promethazine, *in vitro* inhibitors of aldehyde oxidase, did not show relevant effects on the formation of penciclovir *in vivo* [see *Drug Interactions* (7.2)].

**Elimination:** Approximately 94% of administered radioactivity was recovered in urine over 24 hours (83% of the dose was excreted in the first 6 hours) after the administration of 5 mg/kg radiolabeled penciclovir as a 1-hour infusion to 3 healthy male volunteers. Penciclovir accounted for 91% of the radioactivity excreted in the urine.

Following the oral administration of a single 500 mg dose of radiolabeled famciclovir to 3 healthy male volunteers, 73% and 27% of administered radioactivity were recovered in urine and feces over 72 hours, respectively. Penciclovir accounted for 82% and 6-deoxy penciclovir accounted for 7% of the radioactivity excreted in the urine. Approximately 60% of the administered radiolabeled dose was collected in urine in the first 6 hours.

After intravenous administration of penciclovir in 48 healthy male volunteers, mean ± SD total plasma clearance of penciclovir was 36.6±3 L/hr (0.48±0.09 L/hr/kg). Penciclovir renal clearance accounted for 74.5±8.8% of total plasma clearance.

Renal clearance of penciclovir following the oral administration of a single 500 mg dose of famciclovir to 109 healthy male volunteers

was 27.7±7.6 L/hr. Active tubular secretion contributes to the renal elimination of penciclovir.

The plasma elimination half-life of penciclovir was 2.0±0.3 hours after intravenous administration of penciclovir to 48 healthy male volunteers and 2.3±0.4 hours after oral administration of 500 mg famciclovir to 124 healthy male volunteers. The half-life in 17 patients with herpes zoster was 2.8±1.0 hours and 2.7±1.0 hours after single and repeated doses, respectively.

#### Special populations:

**Geriatric patients:** Based on cross study comparison, penciclovir AUC was 40% higher and penciclovir renal clearance was 22% lower in elderly subjects (n=18, age 65 to 79 years) as compared with younger subjects. Some of this difference may be due to differences in renal function between the 2 groups. No famciclovir dosage adjustment based on age is recommended unless renal function is impaired [see *Dosage and Administration* (2.3)]. *Use in Specific Populations* (8.5.1).

**Patients with renal impairment:** In subjects with varying degrees of renal impairment, apparent plasma clearance, renal clearance, and the plasma-elimination rate constant of penciclovir decreased linearly with reductions in renal function, after both single and repeated dosing [see *Use in Specific Populations* (8.6)]. A dosage adjustment is recommended for patients with renal impairment [see *Dosage and Administration* (2.3)].

**Patients with hepatic impairment:** Mild or moderate hepatic impairment had no effect on the extent of availability (AUC) of penciclovir [see *Use in Specific Populations* (8.7)]. No dosage adjustment is recommended for patients with mild or moderate hepatic impairment. The effect of severe hepatic impairment on the pharmacokinetics of penciclovir has not been evaluated.

**HIV-infected patients:** Following oral administration of a single dose of 500 mg famciclovir to HIV-positive patients, the pharmacokinetic parameters of penciclovir were comparable to those observed in healthy subjects.

**Gender:** The pharmacokinetics of penciclovir were evaluated in 18 healthy male and 18 healthy female volunteers after single-dose oral administration of 500 mg famciclovir. AUC of penciclovir was 9.3±1.9 mcg hr/mL and 11.1±2.1 mcg hr/mL in males and females, respectively. Penciclovir renal clearance was 28.5±8.9 L/hr and 21.8±4.3 L/hr, respectively. These differences were attributed to differences in renal function between the 2 groups. No famciclovir dosage adjustment based on gender is recommended.

**Race:** A retrospective evaluation was performed to compare the pharmacokinetic parameters obtained in black and Caucasian subjects after single and repeat once-daily, twice-daily, or three times-daily administration of famciclovir 500 mg. Data from a study in healthy volunteers (single dose), a study in subjects with varying degrees of renal impairment (single and repeat dose) and a study in subjects with hepatic impairment (single dose) did not indicate any significant differences in the pharmacokinetics of penciclovir between black and Caucasian subjects.

#### 12.4 Microbiology

**Mechanism of action:** Famciclovir is a prodrug of penciclovir, which has demonstrated inhibitory activity against herpes simplex virus types 1 (HSV-1) and 2 (HSV-2) and varicella zoster virus (VZV). In cells infected with HSV-1, HSV-2 or VZV, the viral thymidine kinase phosphorylates penciclovir to a monophosphate form that, in turn, is converted by cellular kinases to the active form penciclovir triphosphate. Biochemical studies demonstrate that penciclovir triphosphate inhibits HSV-2 DNA polymerase competitively with deoxyguanosine triphosphate. Consequently, α-herpes viral DNA synthesis and, therefore, replication are selectively inhibited. Penciclovir triphosphate has an intracellular half-life of 10 hours in HSV-1, 20 hours in HSV-2, and 7 hours in HSV-2-infected cells grown in culture. However, the clinical significance of the intracellular half-life is unknown.

**Antiviral activity:** In cell culture studies, penciclovir has antiviral activity against the following herpes viruses: HSV-1, HSV-2 and VZV. The antiviral activity of penciclovir against wild type strains grown on human foreskin fibroblasts was assessed with a plaque reduction assay and staining with crystal violet 3 days post-infection for HSV and 10 days post-infection for VZV. The median EC<sub>50</sub> values of penciclovir against laboratory and clinical isolates of HSV-1, HSV-2 and VZV were 2 µM (range 1.2 to 2.4 µM, n=7), 2.6 µM (range 1.6 to 11 µM, n=6), and 34 µM (range 6.7 to 17 µM, n=6), respectively.

#### Resistance

##### In Cell Culture

Penciclovir-resistant HSV-1 and HSV-2 strains were isolated in cell culture. Penciclovir-resistant mutants of HSV and VZV resulted from mutations in the viral thymidine kinase (TK) and DNA polymerase (POL) genes. Frameshifts were commonly isolated and result in premature truncation of the HSV TK product with decreased enzymatic activity and consequent decreased susceptibility to penciclovir. Mutations in the viral TK gene may lead to complete loss of TK activity (TK negative), reduced levels of TK activity (TK partial), or alteration in the ability of viral TK to phosphorylate the drug without an equivalent loss in the ability to phosphorylate thymidine (TK altered). In cell culture, the following resistance-associated substitutions in TK of HSV-1 and HSV-2 were observed: HSV-1 TK G6C, F13L, H142Y, G200D, L205S, S254Stop, V267G, and T287M; HSV-2 TK G39E, F191L, E226K, and T288M. The median EC<sub>50</sub> values observed in a plaque reduction assay with penciclovir resistant HSV-1, HSV-2, and VZV were 69 µM (range 14 to 115 µM, n=6), 46 µM (range 4 to > 395 µM, n=9), and 92 µM (range 51 to 148 µM, n=4), respectively.

Resistance and Cross-resistance in Clinical Isolates from HSV-Infected Patients

Clinical HSV-1 and HSV-2 isolates obtained from patients who failed treatment with acyclovir for their α-herpesvirus infections were evaluated for genotypic changes in the TK and POL genes. These HSV isolates had frameshift mutations leading to loss of thymidine kinase or had substitutions in the viral thymidine kinase and viral DNA polymerase. Phenotypic analysis of these clinical isolates confirmed resistance to penciclovir and acyclovir. These and other resistance-associated substitutions reported in the literature, or observed in clinical trials, are listed in Table 6. The list is not all inclusive and additional changes will likely be identified in HSV variants isolated from patients who fail penciclovir containing regimens. The possibility of viral resistance to penciclovir should be considered in patients who fail to respond or experience recurrent viral shedding during therapy.

**Table 6: Summary of Known HSV TK and POL Amino Acid Substitutions Confering Resistance to Acyclovir and Cross-Resistance to Penciclovir**

| HSV-1 | TK  | G6C, R32H, R51W, Y53C/H, H58N, G59W, G61A, S74SStop, E83K, P84L, T103P, Q104Stop, D116N, M121R, I143V, R163H, L170P, Y172C, A174P, R1760/W, Q185R, A189V, G200D, G206R, L208H, R216C, R220H, R222C/H, FS 224, Y239S, T245M, Q250SStop, S254Stop, R256W, Q261Stop, R281Stop, T287M, L315S, M322K, C336Y |
|-------|-----|--|
| HSV-2 | TK  | G39E, R51W, Y53N, R177W*, R221H, T288M*  |
| HSV-1 | POL | A657T, D672N, V715G, A719V, S724N, E798K, V813M, N815S, Y818C, G841S, R842S, F891C, V958L  |
| HSV-2 | POL | -  |

\*These substitutions were also observed in penciclovir-treated patients.

Note: Many additional pathways to penciclovir resistance likely exist.

Cross-resistance has been observed among HSV isolates carrying foscarnet resistance-associated substitutions (Table 7).

**Table 7: Summary of Known HSV-1 POL Amino Acid Substitutions Confering Resistance to Foscarnet and Cross-Resistance to Penciclovir**

| HSV-1 | POL | D672N, S724N, E798K, V813M, Y818C, F891C, V958L |
|-------|-----|---|
|-------|-----|---|

#### 13 NONCLINICAL TOXICOLOGY

##### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**Carcinogenesis:** Two-year dietary carcinogenicity studies with famciclovir were conducted in rats and mice. An increase in the incidence of mammary adenocarcinoma (a common tumor in animals of this strain) was seen in female rats receiving the high dose of 600 mg/kg/day (1.1 to 4.5x the human systemic exposure at the recommended total daily oral dose ranging between 500 mg and 2000 mg, based on area under the plasma concentration curve comparisons [24 hr AUC] for penciclovir). No increases in tumor incidence were reported in male rats treated at doses up to 240 mg/kg/day (0.7 to 2.7x the human AUC), or in male and female mice at doses up to 600 mg/kg/day (0.3 to 1.2x the human AUC).

**Mutagenesis:** Famciclovir and penciclovir (the active metabolite of famciclovir) were tested for genotoxic potential in a battery of *in vitro* and *in vivo* assays. Famciclovir and penciclovir were negative in *in vitro* tests for gene mutations in bacteria (*S. typhimurium* and *E. coli*) and unscheduled DNA synthesis in mammalian HaLa 83 cells (at doses up to 10,000 and 5,000 mcg/plate, respectively). Famciclovir was also negative in the L5178Y mouse lymphoma assay (5000 mcg/mL), the *in vivo* mouse micronucleus test (4800 mg/kg), and rat dominant lethal study (5000 mg/kg). Famciclovir induced increases in polyploidy in human lymphocytes *in vitro* in the absence of chromosomal damage (1200 mcg/mL). Penciclovir was positive in the L5178Y mouse lymphoma assay for gene mutation/chromosomal aberrations, with and without metabolic activation (1000 mcg/mL) in human lymphocytes; penciclovir caused chromosomal aberrations in the absence of metabolic activation (250 mcg/mL). Penciclovir caused an increased incidence of micronuclei in mouse bone marrow *in vivo* when administered intravenously at doses highly toxic to bone marrow (500 mg/kg), but not when administered orally.

**Impairment of fertility:** Testicular toxicity was observed in rats, mice, and dogs following repeated administration of famciclovir or penciclovir. Testicular changes included atrophy of the seminiferous tubules, reduction in sperm count, and/or increased incidence of sperm with abnormal morphology or reduced motility. The degree of toxicity to male reproduction was related to dose and duration of exposure. In male rats, decreased fertility was observed after 10 weeks of dosing at 500 mg/kg/day (1.4 to 5.7x the human AUC).

The no observable effect level for sperm and testicular toxicity in rats following chronic administration (26 weeks) was 50 mg/kg/day (0.15 to 0.6x the human systemic exposure based on AUC comparisons). Testicular toxicity was observed following chronic administration to mice (104 weeks) and dogs (26 weeks) at doses of 600 mg/kg/day (0.3 to 1.2x the human AUC) and 150 mg/kg/day (1.3 to 5.1x the human AUC), respectively.

Famciclovir had no effect on general reproductive performance or fertility in female rats at doses up to 1000 mg/kg/day (2.7 to 10.8x the human AUC).

#### 14 CLINICAL STUDIES

##### 14.1 Herpes Labialis (Cold Sores)

A randomized, double-blind, placebo-controlled trial was conducted in 701 immunocompetent adults with recurrent herpes labialis. Patients self-initiated therapy within 1 hour of first onset of signs or symptoms of a recurrent herpes labialis episode with famciclovir 1500 mg as a single dose (n=227), famciclovir 750 mg twice daily (n=220) or placebo (n=254) for 1 day. The median time to healing among patients with non-aborted lesions (progressing beyond the papule stage) was 4.4 days in the famciclovir 1500 mg single-dose group (n=152) as compared to 6.2 days in the placebo group (n=168). The median difference in time to healing between the placebo and famciclovir 1500 mg treated groups was 1.3 days (95% CI: 0.6-2.0). No differences in proportion of patients with aborted lesions (not progressing beyond the papule stage) were observed between patients receiving famciclovir or placebo: 33% for famciclovir 1500 mg single dose and 34% for placebo. The median time to loss of pain and tenderness was 1.7 days in famciclovir 1500 mg single dose-treated patients vs. 2.9 days in placebo-treated patients.

##### 14.2 Genital Herpes

###### Recurrent episodes:

A randomized, double-blind, placebo-controlled trial was conducted in 329 immunocompetent adults with recurrent genital herpes. Patients self-initiated therapy within 6 hours of the first sign or symptom of a recurrent genital herpes episode with either famciclovir 1000 mg twice daily (n=163) or placebo (n=166) for 1 day. The median time to healing among patients with non-aborted lesions (progressing beyond the papule stage) was 4.3 days in famciclovir-treated patients (n=125) as compared to 6.1 days in placebo-treated patients (n=145). The median difference in time to healing between the placebo and famciclovir-treated groups was 1.2 days (95% CI: 0.5 to 2.0). Twenty-three percent of famciclovir-treated patients had aborted lesions (no lesion development beyond erythema) vs. 13% in placebo-treated patients. The median time to loss of all symptoms (e.g., tingling, itching, burning, pain, or tenderness) was 3.3 days in famciclovir-treated patients vs. 5.4 days in placebo-treated patients. A randomized (2,1), double-blind, placebo-controlled trial was conducted in 304 immunocompetent black and African American adults with recurrent genital herpes. Patients self-initiated therapy within 6 hours of the first sign or symptom of a recurrent genital herpes episode with either famciclovir 1000 mg twice daily (n=206) or placebo (n=98) for 1 day. The median time to healing among patients with non-aborted lesions was 5.4 days in famciclovir-treated patients (n=152) as compared to 4.8 days in placebo-treated patients (n=78). The median difference in time to healing between the placebo and famciclovir-treated groups was -0.26 days (95% CI: -0.98 to 0.40).

**Suppressive therapy:** Two randomized, double-blind, placebo-controlled, 12-month trials were conducted in 934 immunocompetent adults with a history of 6 or more recurrences of genital herpes episodes per year. Comparisons included famciclovir 125 mg three times daily, 250 mg twice daily, 250 mg three times daily, and placebo. At 12 months, 60% to 65% of patients were still receiving famciclovir and 25% were receiving placebo treatment. Recurrence rates at 6 and 12 months in patients treated with the 250 mg twice daily dose are shown in Table 8.

**Table 8: Recurrence Rates at 6 and 12 Months in Adults with Recurrent Genital Herpes on Suppressive Therapy**

|   | Recurrence Rates at 6 Months           |                 | Recurrence Rates at 12 Months          |                 |
|---|--|-----------------|--|-----------------|
|   | Famciclovir 250 mg twice daily (n=236) | Placebo (n=233) | Famciclovir 250 mg twice daily (n=236) | Placebo (n=233) |
| Recurrence-free                             | 39%                                    | 10%             | 29%                                    | 6%              |
| Recurrences <sup>1</sup>                    | 47%                                    | 74%             | 53%                                    | 78%             |
| <sup>2</sup> Lost to follow-up <sup>3</sup> | 14%                                    | 16%             | 17%                                    | 16%             |

<sup>1</sup>Based on patient reported data; not necessarily confirmed by a physician.

<sup>2</sup>Patients recurrence-free at time of last contact prior to withdrawal.

Famciclovir-treated patients had approximately one-fifth the median number of recurrences as compared to placebo-treated patients. Higher doses of famciclovir were not associated with an increase in efficacy.

##### 14.3 Recurrent Orolabial or Genital Herpes in HIV-Infected Patients

A randomized, double-blind trial compared famciclovir 500 mg twice daily for 7 days (n=150) with oral acyclovir 400 mg 5 times daily for 7 days (n=143) in HIV-infected patients with recurrent orolabial or genital herpes treated within 48 hours of lesion onset. Approximately 40% of patients had a CD<sub>4</sub> count below 200 cells/mm<sup>3</sup>. 54% of patients had anogenital lesions and 36% had orolabial lesions. Famciclovir therapy was comparable to oral acyclovir in reducing new lesion formation and in time to complete healing.

##### 14.4 Herpes Zoster (Shingles)

Two randomized, double-blind trials, 1 placebo-controlled and 1 active-controlled, were conducted in 964 immunocompetent adults with reconstituted herpes zoster. Treatment was initiated within 72 hours of first lesion appearance and was continued for 7 days.

In the placebo-controlled trial, 419 patients were treated with either famciclovir 500 mg three times daily (n=138), famciclovir 750 mg three times daily (n=135) or placebo (n=146). The median time to full crusting was 5 days among famciclovir 500 mg-treated patients as compared to 7 days in placebo-treated patients. The times to full crusting, loss of vesicles, loss of ulcers, and loss of crusts were shorter for famciclovir 500 mg-treated patients than for placebo-treated patients in the overall study population. The effects of famciclovir were greater when therapy was initiated within 48 hours of rash onset; it was also more profound in patients 50 years of age or older. Among the 65.2% of patients with at least 1 positive viral culture, famciclovir-treated patients had a shorter median duration of viral shedding than placebo-treated patients (1 day and 2 days, respectively).

There were no overall differences in the duration of pain before rash healing between famciclovir- and placebo-treated groups. In addition, there was no difference in the incidence of pain after rash healing (postherpetic neuralgia) between the treatment groups. In the 186 patients (44.4% of total study population) who developed postherpetic neuralgia, the median duration of postherpetic neuralgia was shorter in patients treated with famciclovir 500 mg than in those treated with placebo (63 days and 119 days, respectively). No additional efficacy was demonstrated with higher dose of famciclovir.

In the active-controlled trial, 545 patients were treated with 1 of 3 doses of famciclovir three times daily or with acyclovir 800 mg five times daily. Times to full lesion crusting and times to loss of acute pain were comparable for all groups and there were no statistically significant differences in the time to loss of postherpetic neuralgia between famciclovir and acyclovir-treated groups.

##### 16 HOW SUPPLIED/STORAGE AND HANDLING

Famciclovir tablets are supplied as film-coated tablets as follows: 125 mg in bottles of 30; 250 mg in bottles of 30; and 500 mg in bottles of 30.

Famciclovir 125 mg tablet:

White coloured, circular, film-coated, biconvex tablet debossed with 'C' and '269' on one side and plain on other side.

125 mg 30's ..... NDC 69097-269-02